

Missouri School Immunization Requirements Screening Tool

*This chart is a **basic screening tool** for use in evaluating the immunization status of students enrolling in grades Kindergarten, Eighth and Twelfth. There may be some additional spacing requirements not included on this **basic screening tool**.*

- STEP 1:** Review the immunization requirements for the student's grade.
- STEP 2:** Count the number of doses required for each vaccine category.
- STEP 3:** Check dose and spacing on DTaP, Tdap, MCV, Polio, MMR, Hepatitis B, Varicella.
- STEP 4:** If an In progress card is on file, check the due date for the next dose. Due to the spacing requirements of the vaccine series, this appointment must be kept. If the appointment is **not** kept the child is no longer in progress and is noncompliant.
- STEP 5:** If student has a Religious Immunization Exemption (Imm.P.11A) or Medical Immunization Exemption (Imm.P.12) the appropriate exemption card must be on file.

VACCINES REQUIRED	DOSES REQUIRED BY GRADE			DOSE/SPACING REQUIREMENTS If the vaccine is given 4 days early, the child is considered adequately immunized.
	K	8 th	12 th	
DTaP/DT	4+	4+	4+	Last dose on or after the fourth birthday and the last dose of pediatric pertussis before the seventh birthday. Maximum needed: six doses.
Tdap		1	1	Tdap, which contains pertussis vaccine, is required. If a student received a Tdap, the student is up-to-date. Tdap is currently licensed for one dose only; an additional dose is not needed.
MCV		1	2	<u>Grade 8:</u> One dose of MCV is required. • Must be given at age 10 or older. <u>Grade 12:</u> Two doses of MCV is required unless the first dose was administered to a student who was 16 years of age or older, in which case only one dose is required. Note: Meningococcal B vaccine will not count for the MCV requirement.
IPV (Polio)	3+	3+	3+	<u>Kindergarten-6 Grade:</u> Last dose must be administered on or after the fourth birthday. The interval between the next-to-last and last dose should be at least six months. <u>7-12 Grades:</u> Last dose on or after the fourth birthday. If all four doses are administered appropriately and received prior to the fourth birthday, an additional dose is not needed. Any combination of four doses of IPV and OPV by four-six years of age constitutes a complete series. Maximum needed: four doses.
MMR	2	2	2	First dose must be given on or after twelve months of age.
Hepatitis B	3+	3+	3+	To be considered valid the following must be met: • At least four weeks between dose one and two • At least 8 weeks between doses two and three • At least 16 weeks between doses one and three • Final dose must be at given at least 24 weeks of age.
Varicella	2	1		First dose must be given on or after twelve months of age.

